

AD-A092 536

MASSACHUSETTS UNIV AMHERST DEPT OF MATHEMATICS AND S--ETC F/6 12/1
A CHARACTERIZATION OF A POLYA-EGGENBERGER AND OTHER DISCRETE DI--ETC(U)
SEP 80 R M KORWAR AFOSR-80-0219

UNCLASSIFIED

TR-8

AFOSR-TR-80-1180

NL

1 of 1

40
900508



END

DATE
FILMED

1-81
DTIC

AD A092536

| REPORT DOCUMENTATION | | READ INSTRUCTIONS BEFORE COMPLETING FORM |
|--|--|---|
| 1. REPORT NUMBER | 2. AUTHOR | 3. RECIPIENT'S CATALOG NUMBER |
| AFOSR-TR- 30 - 1180 | AD-A092536 | 536 |
| 4. TITLE (and Subtitle) | 5. TYPE OF REPORT & PERIOD COVERED | |
| A CHARACTERIZATION OF A POLYA EGGENBERGER AND OTHER DISCRETE DISTRIBUTIONS BY RECORD VALUES | Interim | |
| 7. AUTHOR(s) | 6. PERFORMING ORG. REPORT NUMBER | |
| Ramesh M. Korwar | | |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS | 8. CONTRACT OR GRANT NUMBER(s) | |
| The University of Massachusetts Department of Mathematics and Statistics Amherst, Massachusetts 01003 | AFOSR 80-0219 <i>new</i> | |
| 11. CONTROLLING OFFICE NAME AND ADDRESS | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS | |
| Air Force Office of Scientific Research /NM Bolling AFB, Washington, DC 20332 | 61102F 2304/A5 | |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) | 12. REPORT DATE | |
| | September 1980 | |
| | 13. NUMBER OF PAGES | |
| | 8 | |
| | 15. SECURITY CLASS. (of this report) | |
| | UNCLASSIFIED | |
| | 15a. DECLASSIFICATION DOWNGRADING SCHEDULE | |
| 16. DISTRIBUTION STATEMENT (of this Report) | | |
| Approved for public release; distribution unlimited | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | |
| DTIC ELECTE DEC 05 1980 | | |
| 18. SUPPLEMENTARY NOTES | | |
| E | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) | | |
| Characterization Linear regression Discrete distributions Polya Eggenberger Generalized hypergeometric Record Values | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) | | |
| <i>note: all n - print "sub"</i> Let X_1, X_2, \dots , be a sequence of independent and identically distributed discrete random variables. Define the sequence $N(n)$ by $N(1)=1, N(n)=\min \{j j N(n-1), X_j X_N(n-1)\}, n=2,3, \dots$. Let $R_n = X_{N(n)}$. Then R_n is the sequence of record values. By convention $R_1 = X_1$. Here a characterization of a Polya Eggenberger and other discrete distributions including the geometric, is made by the linearity of regression of $R_2 - R_1$ on R_1 . | | |

DDC FILE COPY

(18) AFOSR-TR-80-1180

(11) 2384

(12) 91

(6)

A CHARACTERIZATION OF A
POLYA-EGGENBERGER AND OTHER DISCRETE DISTRIBUTIONS
BY RECORD VALUES.

by

(10) Ramesh M. Korwar*

(16) 2384

(9)
AFOSR Technical Report No. 8

(17) A51

(14) 11

(15) ✓ AFOSR-TR-80-1180

| | |
|--------------------|--|
| Accession For | |
| NTIS GRA&I | <input checked="checked" type="checkbox"/> |
| DDC TAB | <input type="checkbox"/> |
| Unannounced | <input type="checkbox"/> |
| Justification | |
| By | |
| Distribution/ | |
| Availability Codes | |
| Dist | or |
| A | al |

September, 1980

The University of Massachusetts
Department of Mathematics and Statistics
Amherst, Massachusetts

Research sponsored by the Air Force Office of Scientific Research, AFOSR, USAF,
under Grant AFOSR-80-0219.

80 12 01 019

Approved for release
Distribution unlimited.

4-2-377

A Characterization of a Polya-Eggerberger and Other
Discrete Distributions by Record Values

Ramesh M. Korwar

University of Massachusetts

Abstract

Let X_1, X_2, \dots , be a sequence of independent and identically distributed discrete random variables. Define the sequence $W(n)$ by $W(1)=1, W(n)=\min\{j \mid X_j > X_{j-1}, X_j > X_{j-2}, \dots, X_j > X_1\}$, $n=2, \dots$. Let $R_n = X_{W(n)}$. Then R_n is the sequence of record values. By convention $R_1 = X_1$. Here a characterization of a Polya-Eggerberger and other discrete distributions including the geometric, is made by the linearity of regression of $R_2 - R_1$ on R_1 .

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
MONTGOMERY, ALABAMA 36115
This report has been reviewed and is
approved for release under E.O. 12812 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

A CHARACTERIZATION OF A
POLYA-EGGENBERGER AND OTHER DISCRETE
DISTRIBUTIONS BY RECORD VALUES ¹

by

R M KORWAR
Indian Statistical Institute
and
University of Massachusetts

SUMMARY. Let X_1, X_2, \dots , be a sequence of independent and identically distributed discrete random variables. Define the sequence $\{N(n)\}$ by $N(1)=1, N(n)=\min\{j \mid j > N(n-1), X_j > X_{N(n-1)}\}$, $n=2, 3, \dots$. Let $R_n = X_{N(n)}$. Then $\{R_n\}$ is the sequence of record values. By convention $R_1 = X_1$. Here a characterization of a Polya-Eggenberger and other discrete distributions including the geometric, is made by the linearity of regression of $R_2 - R_1$ on R_1 .

Consider a sequence X, X_1, X_2, \dots of independent and identically distributed (i.i.d.) discrete random variables (r.v.) with

$$P(X=j) = p_j, j=0, \dots, m \quad (1)$$

Here m is either a positive integer or ∞ . Define

$$N(1)=1, N(n)=\min\{j \mid j > N(n-1), X_j > X_{N(n-1)}\}, n=2, 3, \dots,$$

$R_1 = X_1$ and $R_n = X_{N(n)}$, $n=2, 3, \dots$. Then $\{R_n\}$ is the sequence of record values and $\{N(n)\}$ the sequence of times at which record values occur.

In this note we characterize the geometric, a Polya-Eggenberger and a generalized hypergeometric distribution by linearity of regression of $R_2 - R_1$ on R_1 . Thus the

¹Research sponsored by the Air Force Office of Scientific Research, AFOSR, USAF, under Grant AFOSR-60-0219.

characterization given here extends Theorem 2 of Srivastava (1979) characterizing the geometric by the constancy of regression of $R_2 - R_1$ on R_1 .

Consider for X one of the following distributions.

$$p_j = (1-p_0) c \{1-c\}^{j-1}, \quad j=1,2,\dots, \quad (2)$$

$0 < c < 1.$

$$p_j = (1-p_0) \binom{a}{j-1} \binom{b}{n-j+1} / \binom{a+b}{n}, \quad j=1,\dots, n+1, \quad (3)$$

$a < 0; n > 0; n \text{ integral}; n-b-1 > 0; b < 0, b \neq -1.$

and

$$p_j = (1-p_0) \binom{a}{j-1} \binom{b}{n-j+1} / \binom{a+b}{n}, \quad j=1,2,\dots, \quad (3a)$$

$a < 0; n < 0; n-b-1 < 0; a+b+1 > 0.$

Distribution (2) is a geometric distribution, (3) and (3a) are versions of generalized hypergeometric distribution; Type 2A and Type IV, as described by Kemp and Kemp (1956). For $a=-1$, (3) can be recast as

$$p_j = (1-p_0) \binom{n}{j-1} \frac{A(A+s), \dots, (A+j-2s) \quad B(B+s), \dots, (B+n-(j-1)-1s)}{(A+B) (A+B+s) \dots (A+B+n-1s)} \quad (3b)$$

$j=1,\dots, n+1$

with $A=1$, $B=-b$, $s=1$. An example of a Polya-Eggenberger distribution is (3b).

We are now ready to prove the

Theorem: Suppose X takes on only the values $0, \dots, n$ with positive probabilities and assume that X has a finite expectation. Then

$$E(R_2 - R_1 \mid R_1 = i) = \alpha + i\beta \quad \text{a.s.}$$

$$\alpha, \beta \text{ Constants} \quad (4)$$

$$i = 0, \dots, m-1$$

if, and only if, X has either the geometric (2), or a Polya-Eggenberger (3b) with $A=1$ and $s=1$, or a generalized hypergeometric distribution (3a) with $a=-1$. Furthermore, $\beta=0$ iff X is geometric (2), $\beta < 0$, $\beta \neq 1$ iff X is (3b) with $A=1$, $s=1$ and $\beta > 0$ iff X is (3a) with $a=-1$.

Proof: First consider the 'only if' part. Suppose (4) holds. Now it is shown in Srivastava (1979) that

$$E(R_2 - R_1 \mid R_1 = i) = \frac{\sum_{j=1}^{m-i} p_{i+j}}{\sum_{j=i+1}^m p_j}, \quad i=0, \dots, m-1$$

Thus it follows that

$$\sum_{j=1}^{m-i} j p_{i+j} = (\alpha + i\beta) \sum_{j=i+1}^m p_j \quad (5)$$

and

$$\sum_{j=1}^{m-i-1} j p_{i+j+1} = (\alpha + i\beta) \sum_{j=i+2}^m p_j \quad (6)$$

Subtracting (6) from (5) we have

$$(1+\beta) \sum_{j=i+1}^m p_j = (\alpha + i\beta) p_{i+1} \quad (7)$$

Replacing i by $(i+1)$ in (7) and subtracting the result from (7), we finally obtain

$$p_{i+2} / p_{i+1} = (\alpha + \beta i - 1) / \{ \alpha + \beta(i+2) \}, \quad i=0, 1, \dots, m-2.$$

This last result and (7) yield

$$p_j = p_1 \frac{(\alpha-1) \dots (\alpha+j-2\beta-1)}{(\alpha+2\beta) \dots (\alpha+j\beta)} \quad j=1, \dots, m. \quad (8)$$

and

$$p_1 = (1+\beta) (1-p_0) / (\alpha+\beta). \quad (9)$$

From (4) it follows that

$$\alpha \geq 1 \quad (10)$$

We consider three cases (i) $\beta=0$, (ii) $\beta<0$ and (iii) $\beta>0$. The case $\beta=0$ is already considered by Srivastava (1979). His result is included in (8) and specializing for $\beta=0$, (8) and (9) give (2) with $c=1/\alpha$.

Next consider the cases $\beta \neq 0$. Now, from (4), it follows that

$$\beta < 0 \quad \text{if, and only if, } X \text{ is bounded} \quad (11)$$

$$\text{If } \beta < 0 \quad \text{then } \alpha + \beta(m-1) = 1 \quad (12)$$

We obtain (12) from (4) by specializing $i=m-1$. From (8) and (9) it is clear that $\beta=-1$ leads to the degeneracy of X at 0, a contradiction of the assumption of m being a positive integer. For the cases $\beta \neq 0$ (8) can be recast as

$$p_j = p_1 \left\{ \frac{A^{[j-1]} B^{[j-1]}}{C^{[j-1]} (j-1)!} \right\}, \quad j=1, \dots, m \quad (13)$$

where x is the ascending factorial,

$$x^{[j]} = x(x+1)\dots(x+j-1), \quad j=1, 2, \dots, \quad x^{[0]} = 1;$$

p_1 is given by (9) and

$$A = (\alpha-1)/\beta \quad B=1, \quad \text{and } C = \alpha/\beta+2. \quad (14)$$

The product in the curled brackets in (13) is the coefficient of z^{j-1} in the hypergeometric function

$$F(A;B;C;z) = \sum_{j=0}^{\infty} \left\{ \frac{[j]_A [j]_B}{[j]_C j!} \right\} z^j.$$

Thus (13) is the generalized hypergeometric distribution (3) (or 3(a)) with

$$a=-1, b=1+1/\beta, n=-(\alpha-1)/\beta \quad (15)$$

First suppose $\beta < 0$. Then, from (12) and (15) we have that $n=m-1$, and if $m \geq 2$, from (9), that $b < 0$. Obviously, $n-b-1 > 0$ for $m \geq 2$. Thus, for the case $\beta < 0$, $\beta \neq 1$ and $m \geq 2$ we have a generalized hypergeometric distribution (3) with parameters $a=-1$, $b=1+1/\beta$ and $n=m-1$. This distribution is also a Polya-Eggenberge distribution (3b) with parameter $A=1$, $B=-1/\beta$, $n=m-1$ and $s=1$. However, (4) will not lead to any particular distribution if $m=1$.

Finally assume $\beta > 0$. From (15), we have that X has the generalized hypergeometric distribution (3a) with the parameters given by (15).

Now for the "if" part. Let X have (3) or (3a) with $a=-1$. Then we have

$$p_{j+1}/p_j = (j-1-n) / (b-n+j), \quad j=1,2,\dots,m-1.$$

from which we get

$$\sum_{j=k+1}^m j p_j / \sum_{j=k+1}^m p_j = (b-n-1+bk) / (b-1), \quad k=0,\dots,m-1$$

-o 6 o-

which in turn yields (4) with $\alpha=(b-n-1)/(b-1)$ and $\beta=b/(b-1)$. Hence the 'if' part is proved.

REFERENCES

- KEMP, C.D. and KEMP, A.W. (1956): Generalized Hypergeometric Distributions. J. Roy. Stat. Soc. Ser B, 18, 202-11.
- SRIVASTAVA, R.C. (1979): Two Characterizations of the Geometric Distribution By Record Values. Sankhyā Ser. B, 40, 276-8.

DATE
FILMED
- 8